arXiv:1508.06930v1 [math.CO] 27 Aug 2015

LATTICE PATHS, YOUNG TABLEAUX, AND WEIGHT MULTIPLICITIES

REBECCA L. JAYNE AND KAILASH C. MISRA

Abstract. For $\ell \geq 1$ and $k \geq 2$, we consider certain admissible sequences of $k-1$ lattice paths in a colored $\ell \times \ell$ square. We show that the number of such admissible sequences of lattice paths is given by the sum of squares of the number of standard Young tableaux of partitions of ℓ with height $\leq k$, which is also the number of $(k+1)k\cdots 21$-avoiding permutations of $\{1, 2, \ldots, \ell\}$. Finally, we apply this result to the representation theory of the affine Lie algebra $\widehat{sl}(n)$ and show that this quantity gives the multiplicity of certain maximal dominant weights in the irreducible module $V(k\Lambda_0)$.

1. Introduction

For fixed integers $\ell \geq 1$ and $k \geq 2$, we consider the admissible sequences of $k-1$ lattice paths in a colored $\ell \times \ell$ square given in [1]. Each admissible sequence of paths can be associated with a partition λ of ℓ. In Section 2, we show that the number of self-conjugate admissible sequences of paths associated with λ is equal to the number of standard Young tableaux of shape λ, and thus can be calculated using the hook length formula. We extend this result to include the non-self-conjugate admissible sequences of paths and show that the number of all such admissible sequences of paths is equal to the sum of squares of the number of standard Young tableaux of partitions of ℓ with height less than or equal to k. Using the RSK correspondence in [3], it is shown in ([4], Corollary 7.23.12) that the sum of squares of the number of standard Young tableaux of partitions of ℓ with height less than or equal to k is equal to the number of $(k+1)k\cdots 21$-avoiding permutations of $\{1, 2, \ldots, \ell\}$.

In Section 3, we apply our results to the representation theory of the affine Kac-Moody algebra $\widehat{sl}(n)$. Let $\{\alpha_0, \alpha_1, \ldots, \alpha_{n-1}\}$, $\{h_0, h_1, \ldots, h_{n-1}\}$, and $\{\Lambda_0, \Lambda_1, \ldots, \Lambda_{n-1}\}$ denote the simple roots, simple coroots, and fundamental weights respectively. Note that $\Lambda_j(h_i) = \delta_{ij}$. For $1 \leq \ell \leq \lfloor \frac{n}{2} \rfloor$, set $\gamma_\ell = \ell\alpha_0 + (\ell-1)\alpha_1 + (\ell-2)\alpha_2 + \cdots + \alpha_{\ell-1} + \alpha_{n-\ell+1} + \cdots + (\ell-2)\alpha_{n-2} + (\ell-1)\alpha_{n-1}$ and $\mu_\ell = k\Lambda_0 - \gamma_\ell$. As shown in [1], μ_ℓ are maximal dominant weights of the irreducible $\widehat{sl}(n)$-module $V(k\Lambda_0)$. We show that the multiplicity of the weight μ_ℓ in $V(k\Lambda_0)$ is the number of $(k+1)k\cdots 21$-avoiding permutations of $\{1, 2, \ldots, \ell\}$, which proves Conjecture 4.13 in [1].

2010 *Mathematics Subject Classification.* Primary 05E10, 17B10; Secondary 05A05, 05A17, 17B67.

Key words and phrases. Lattice path and Young tableau and avoiding permutation and affine Lie algebra and weight multiplicity.

KCM: partially supported by Simons Foundation grant # 307555.

2. Lattice Paths

For fixed integers $\ell \geq 1$ and $k \geq 2$, consider the $\ell \times \ell$ square containing ℓ^2 unit boxes in the fourth quadrant so that the top left corner of the square is at the origin. We assign color $a+b$ to a box if its upper left corner has coordinates (a,b). This gives the following $\ell \times \ell$ colored square Y:

0	1	$\cdots$	$\ell-2$	$\ell-1$
-1	0	$\cdots$	$\ell-3$	$\ell-2$
$\vdots$	$\vdots$	$\ddots$	$\vdots$	$\vdots$
$2-\ell$	$3-\ell$	$\cdots$	0	1
$1-\ell$	$2-\ell$	$\cdots$	-1	0

Figure 1. Y

A lattice path p on Y is a path joining the lower left corner $(0,-\ell)$ to the upper right corner $(\ell, 0)$ moving unit lengths up or right. For two lattice paths p, q on Y we say that $p \leq q$ if the boxes above q are also above p. Now, we draw $k-1$ lattice paths, $p_1, p_2, \ldots, p_{k-1}$, on Y such that $p_1 \leq p_2 \leq \cdots \leq p_{k-1}$. For integers i, j, where $2 \leq i \leq k-1$, $-(\ell-1) \leq j \leq \ell-1$, we define $t_i^j \geq 0$ to be the number of j-colored boxes between p_{i-1} and p_i. We define t_1^j to be the number of j-colored boxes below p_1 and t_0^j to be the number of j-colored boxes above p_{k-1}.

Definition 2.1. [1] *We call a sequence of lattice paths $p_1 \leq p_2 \leq \ldots \leq p_{k-1}$ on Y admissible if it satisfies the following conditions:*

1. *p_1 does not cross the diagonal $y = x - \ell$ and*
2. *for $2 \leq i \leq k-1$, $-(\ell-1) \leq j \leq \ell-1$, we have*

$$t_i^j \leq \min\left\{ t_{i-1}^j, \ell - |j| - t_1^j - \sum_{a=1}^{i-1} t_a^j \right\} \text{ and } \begin{cases} t_i^j \leq t_i^{j-1}, & j > 0 \\ t_i^j \leq t_i^{j+1}, & j < 0. \end{cases}$$

Denote by $\mathcal{T}_\ell^k$ the set of all admissible sequences of $k-1$ paths. Notice that there are ℓ 0-colored boxes in Y and hence for any admissible sequence of paths, $\sum_{i=0}^{k-1} t_i^0 = \ell$. In addition, it follows from Definition 2.1(2) that $t_0^0 \geq t_1^0 \geq \cdots \geq t_{k-1}^0$ for any admissible sequence of paths. Thus, we can and do associate an admissible sequence of paths $p_1 \leq p_2 \leq \ldots \leq p_{k-1}$ on Y with a partition $\lambda = (t_0^0, t_1^0, \ldots, t_{k-1}^0)$ of ℓ. In this case, we say that this admissible sequence of paths is of type λ and often draw λ as a Young diagram.

Example 2.2. Figure 2(a) is an element of $\mathcal{T}_6^4$, where p_1, p_2, and p_3 are shown in Figures 2(b), 2(c), and 2(d), respectively. Notice that this admissible sequence of paths is of type $\lambda = (2, 2, 1, 1)$.


(a) (b) (c) (d)

FIGURE 2. Admissible sequence of paths

For a given partition $\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_k) \vdash \ell$, we let f^λ denote the number of standard Young tableaux of shape λ and let $\lambda' = (\lambda'_1, \lambda'_2, \ldots, \lambda'_m)$ be its conjugate partition. For a Young diagram of type λ, the hook length at a box u in row i and column j is given by $h(u) = \lambda_i + \lambda'_j - i - j + 1$. The hook length formula states that $f^\lambda = \dfrac{\ell!}{\Pi_{u\in\lambda} h(u)}$. We denote the height of a standard Young tableau of shape λ by $ht(\lambda)$.

Example 2.3. For the partition $\lambda = (4, 2, 1) \vdash 7$, we have the Young diagram in Figure 3(a). Using the hook length formula, and moving across the rows and then down in the diagram, we calculate that the number of standard Young tableaux of shape λ is $f^\lambda = \dfrac{7!}{6 \cdot 4 \cdot 2 \cdot 1 \cdot 3 \cdot 1 \cdot 1} = 35$. Figure 3(b) shows one such standard Young tableau of shape λ.



(a) (b)

FIGURE 3. Young diagram and standard Young tableau of shape $(4, 2, 1)$

A lattice path on Y is self-conjugate if it is symmetric about the diagonal from $(0,0)$ to $(\ell,-\ell)$. A sequence of lattice paths is self-conjugate if all lattice paths in the sequence are self-conjugate. Next, we give a relationship between self-conjugate admissible sequences of paths and standard Young tableaux. We denote the set of all self-conjugate admissible sequences of $k-1$ paths on the $\ell\times\ell$ colored square Y by S^k_ℓ. Let F^k_ℓ denote the set of all standard Young tableaux with ℓ boxes and height $\le k$. Then $|F^k_\ell| = \sum\limits_{\lambda\vdash\ell, h(\lambda)\le k} f^\lambda$.

Now, we define a map $\tau\colon F^k_\ell \to S^k_\ell$ as follows. Let $X\in F^k_\ell$. We draw the first ℓ moves of each path, $p_1,p_2,\dots,p_{k-1}$, on the $\ell\times\ell$ colored square Y, giving the portion of each path from $(0,-\ell)$ to the diagonal connecting $(0,0)$ to $(\ell,-\ell)$, move by move. In the i^{th} path p_i, the j^{th} move is up when j is in row 2 through row $(i+1)$ and the j^{th} move is right otherwise. Once the first ℓ moves of every path are drawn, we complete each path by reflecting over the diagonal, obtaining a self-conjugate sequence of paths.

Although the image $\tau(X)$ is a self-conjugate sequence of $k-1$ paths by definition, we still need to show that it is an admissible sequence of paths. Notice that for path p_1 in $\tau(X)$, the m^{th} up move, move v, is represented in column m of row 2 of X. Thus, there are at least m entries in row 1 of X less than or equal to v, which correspond to m right moves prior to the m^{th} up move in p_1 of $\tau(X)$. Hence, p_1 does not cross the diagonal $y = x-\ell$. For $1-\ell\le j\le 0$, we call the line $y=-x+j$ connecting $(0,j)$ and $(\ell+j,-\ell)$ the j-diagonal passing through all j-colored boxes. Note that it takes $\ell+j$ moves to go from $(0,-\ell)$ to the j-diagonal. By definition, t^j_0 is the number of right moves in the path p_{k-1} below the j-diagonal, which equals the number of entries in the first row of X that are less than or equal to $\ell+j$. For $2\le i\le k-1$, the number of up moves in the path p_i below the j-diagonal is equal to the number of j-colored boxes below p_i. Hence, by definition, t^j_i is the number of up moves below the j-diagonal that are in p_i but not in p_{i-1}. Therefore, the number of up moves in p_i (resp. p_{i-1}) below the j diagonal is $t^j_1+t^j_2+\cdots+t^j_i$ (resp. $t^j_1+t^j_2+\cdots+t^j_{i-1}$). Since the paths are self-conjugate, using symmetry it follows that $p_i\ge p_{i-1}$. Hence $p_1\le p_2\le\cdots\le p_{k-1}$. Now observe that by definition of τ, t^j_i equals the number of boxes in row $i+1$ of X with entries less than or equal to $\ell+j$. Since X is a standard tableau, we have $t^j_i\le t^j_{i-1}$. There are at least t^j_1 boxes in the second row of X and hence in the first row of X. So it follows that $2t^j_1+t^j_2+\cdots+t^j_i\le \ell+j$ which implies $t^j_i\le \ell+j-t^j_1-\sum\limits_{a=1}^{i-1}t^j_a$. Hence using the symmetry of the paths we have $t^j_i\le\min\left\{t^j_{i-1},\ell-|j|-t^j_1-\sum\limits_{a=1}^{i-1}t^j_a\right\}$. Finally, for $j<0$, t^j_i (resp. t^{j+1}_i) is the number of boxes in row $i+1$ of X with entries less than or equal to $\ell+j$ (resp. $\ell+j+1$). So $t^j_i\le t^{j+1}_i$ for $j<0$. Now using symmetry and reflecting on the diagonal $y=-x$, we have $t^j_i\le t^{j-1}_i$ for $j>0$, proving that $\tau(X)\in S^k_\ell$.

Theorem 2.4. *For $k\ge 2,\ell\ge 1$, the map $\tau\colon F^k_\ell\to S^k_\ell$ is a bijection. Hence $|S^k_\ell| = \sum\limits_{\lambda\vdash\ell,h(\lambda)\le k} f^\lambda$. In particular, the number of self-conjugate admissible sequences of paths $p_1\le p_2\le\cdots\le p_{k-1}$ of type $\lambda=(t^0_0,t^0_1,\cdots,t^0_{k-1})\vdash\ell$ is equal to the number of standard Young tableaux of shape λ.*

Proof. To prove the statement, it is sufficient to show that the map τ has an inverse. We define the map $\sigma\colon S_\ell^k \to F_\ell^k$ as follows. Let $Z = \{p_1 \leq p_2 \leq \cdots \leq p_{k-1}\} \in S_\ell^k$ be a self-conjugate admissible sequence of paths on the $\ell \times \ell$ square Y of type $\lambda = (t_0^0, t_1^0, \dots, t_{k-1}^0) \vdash \ell$. Note that t_m^0 equals the number of up moves in the path p_m but not in any path $p_i, 1 \leq i < m$. We draw the Young diagram X of shape λ. Now we fill in the boxes of X with numbers $1, 2, \dots, \ell$ as follows. We traverse, in order, the first ℓ moves from $(0, -\ell)$ to the 0-diagonal in each path $p_1, p_2, \dots, p_{k-1}$. If the i^{th} move is a right move for all paths, then we place the number i in the leftmost available box in the first row of X. If the i^{th} move is an up move in any path, then choose m to be the smallest integer such that the up move occurs in the path p_m. In such case, we place the number i in the leftmost available box of row $m+1$ of X.

By construction, the entries in the rows of X increase from left to right. Notice that for $2 \leq i \leq k-1$, any up moves in the path p_i that do not occur in p_{i-1} are represented in row $i+1$ of $X = \sigma(Z)$. So for $-(\ell-1) \leq j \leq 0$, t_i^j in Z is equal to the number of entries in row $(i+1)$ of X that are less than or equal to $\ell + j$. To see that the entries in the columns increase from top to bottom, suppose for the sake of contradiction that for some row r $(2 \leq r \leq k)$ the entry in column c, say a, is less than the entry in column c of row $r-1$. Since the entries in X increase from left to right, the number of entries in row r less than or equal to a is c. Hence $t_{r-1}^{a-\ell} = c$. Since $t_{r-2}^{a-\ell}$ is the number of entries in row $r-1$ less than or equal to a and the entry in column c of row $r-1$ is greater than a, we have $t_{r-2}^{a-\ell} < t_{r-1}^{a-\ell} = c$, which contradicts Definition 2.1. Hence X is a standard Young tableaux and $X \in F_\ell^k$. Using the symmetry of the paths and the definition of τ it follows that $\tau(\sigma(Z)) = Z$, which proves the theorem. ☐

Example 2.5. Notice that Figure 4(a) is an element of F_6^4. To apply τ, we draw three paths to the diagonal, with moves shown in Figure 4(b). Figures 4(c), 4(d), and 4(e) show the first 6 moves of p_1, p_1 and p_2, and p_1, p_2 and p_3, respectively. These paths are reflected over the diagonal to form an element of S_6^4 as shown in Figure 4(f).

Example 2.6. Let $Z \in S_6^4$ as in Figure 5(a). In Figure 5(b), we record the moves for each path. We construct $X \in T_6^4$ as in Figure 5(c) by first drawing the Young diagram of shape $\lambda = (t_0^0 = 2, t_1^0 = 2, t_2^0 = 1, t_3^0 = 1)$. We fill in the integers 1 through 6 using the information collected in Figure 5(b) and applying σ. For example, since move 1 for all paths is right, we place 1 in row 1 of X. Since move 6 has its first up move in p_3, we place 6 in row 4.

Theorem 2.7. *For $k \geq 2$ and $\ell \geq 1$, let $\lambda = (t_0^0, t_1^0, \dots, t_{k-1}^0) \vdash \ell$. Then the number of admissible sequences of $k-1$ paths on a colored $\ell \times \ell$ square Y of type $\lambda = (t_0^0, t_1^0, \dots, t_{k-1}^0)$ is $(f^\lambda)^2$. Hence* $|\mathcal{T}_\ell^k| = \sum_{\lambda \vdash \ell, h(\lambda) \leq k} (f^\lambda)^2$.

Proof. Let $Z_1 = \{p_1 \leq p_2 \leq \cdots \leq p_{k-1}\}$ and $Z_2 = \{p'_1 \leq p'_2 \leq \cdots \leq p'_{k-1}\}$ be two self-conjugate admissible sequences of paths on Y of type $\lambda = (t_0^0, t_1^0, \dots, t_{k-1}^0) \vdash \ell$. Then for $1 \leq i \leq k-1$, combining the portion of the path p_i below the 0-diagonal on Y with the portion of the path p'_i above the 0-diagonal, we obtain an admissible sequence of $k-1$ paths of type λ which is not self-conjugate unless $Z_1 = Z_2$. Conversely, if we have an admissible sequence of paths $Z = \{p_1 \leq p_2 \leq \cdots \leq p_{k-1}\}$ of type $\lambda \vdash \ell$, we can obtain a pair of self-conjugate admissible sequences of paths



(a)

Path	Move 1	2	3	4	5	6
1	R	U	R	R	R	U
2	R	U	R	U	R	U
3	R	U	R	U	U	U

(b)

(c) (d)

(e) (f)

FIGURE 4. Example of τ map

(Z_1, Z_2) each of type λ, where Z_1 (resp. Z_2) is obtained by reflecting the portions of the paths in Z below (resp. above) the 0-diagonal of Y. Thus the set of admissible sequences of $k-1$ paths of type $\lambda \vdash \ell$ is in bijection with the set of pairs of self-conjugate admissible sequences of $k-1$ paths of type $\lambda \vdash \ell$. Hence by Theorem 2.4, the number of admissible sequences of $k-1$ paths in Y of type $\lambda \vdash \ell$ is $(f^\lambda)^2$. Since any partition $\lambda \vdash \ell$ of height less than or equal to k is associated with an admissible sequence of $k-1$ paths of type λ, we have $|\mathcal{T}_\ell^k| = \sum_{\lambda \vdash \ell, h(\lambda) \leq k} (f^\lambda)^2$. $\square$

Example 2.8. In Figure 6(a), we have $Z = \{p_1 \leq p_2 \leq p_3\}$, a non-self-conjugate admissible sequence of paths of type $\lambda = (2, 2, 1, 1)$. Reflecting the portion of Z below the 0-diagonal gives the self-conjugate sequence Z_1 in Figure



(a)

	Move					
Path	1	2	3	4	5	6
1	R	U	R	R	U	R
2	R	U	U	R	U	R
3	R	U	U	R	U	U

(b)

1	4
2	5
3	
6	

(c)

Figure 5. Example of σ map

6(b) and reflecting the portion of Z above the 0-diagonal gives the self-conjugate sequence Z_2 in Figure 6(c), both of type λ. Thus, by the bijection in the proof of Theorem 2.7, the non-self-conjugate sequence of paths Z corresponds to the pair of self-conjugate sequence of paths (Z_1, Z_2).



(a) (b) (c)

Figure 6. Non-self-conjugate admissible sequence of paths correspondence with pair of self-conjugate admissible sequences of paths

3. Weight Multiplicities

In this section we use the result in Theorem 2.7 to prove Conjecture 4.13 in [1], giving a formula for certain weight multiplicities of the affine Lie algebra $\mathfrak{g} = \widehat{sl}(n) = sl(n) \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c \oplus \mathbb{C}d, n \geq 2$, where $sl(n)$ is the simple Lie algebra of $n \times n$ trace zero matrices, c is the canonical central element and $d = 1 \otimes \frac{d}{dt}$ is a degree derivation. We recall some facts about this affine Lie algebra from [2].

The generalized Cartan matrix associated with the affine Lie algebra $\mathfrak{g}$ is the $n \times n$ matrix $A = (a_{ij})_{0 \leq i,j \leq n-1}$ where $a_{ii} = 2, a_{ij} = -1$ for $|i-j| = 1, a_{0,n-1} = a_{n-1,0} = -1$, and $a_{ij} = 0$ otherwise. Let $\Pi = \{\alpha_0, \alpha_1, \ldots, \alpha_{n-1}\}$ and $\Pi^\vee = \{h_0, h_1, \ldots, h_{n-1}\}$ denote the simple roots and simple coroots respectively. We recall that $\alpha_j(h_i) = a_{ij}$ and $\delta = \alpha_0 + \alpha_1 + \cdots + \alpha_{n-1}$ is the null root. The Cartan subalgebra of $\mathfrak{g}$ is $\mathfrak{h} =$span$_\mathbb{C}\{h_1, h_2, \ldots, h_{n-1}, d\}$. The fundamental weights $\{\Lambda_0, \Lambda_1, \ldots, \Lambda_{n-1}\}$ are defined by $\Lambda_i(h_j) = \delta_{ij}, \Lambda_i(d) = 0$ and the set $P = \mathbb{Z}\Lambda_0 \oplus \mathbb{Z}\Lambda_1 \oplus \cdots \oplus \mathbb{Z}\Lambda_{n-1} \oplus \mathbb{Z}\delta$ (resp. $P^\vee = \mathbb{Z}h_0 \oplus \mathbb{Z}h_1 \oplus \cdots \oplus \mathbb{Z}h_{n-1} \oplus \mathbb{Z}d$) is called the weight lattice (resp. coweight lattice).

The set $P^+ = \{\Lambda \in P \mid \Lambda(h_i) \geq 0, \forall i\}$ is called the set of dominant integral weights. For each $\Lambda \in P^+$, there is an irreducible highest weight $\mathfrak{g}$-module $V(\Lambda)$ with highest weight Λ. For $\mu \in \mathfrak{h}^*$, if $V(\Lambda)_\mu = \{v \in V(\Lambda) \mid h(v) = \mu(h)v, \forall h \in \mathfrak{h}\} \neq \{0\}$, then μ is called a weight of $V(\Lambda)$. The multiplicity of μ in $V(\Lambda)$, denoted by $mult_\Lambda(\mu)$, is the dimension of $V(\Lambda)_\mu$. If μ is a weight of $V(\Lambda)$ and $\mu + \delta$ is not a weight, then μ is called a maximal weight. We denote the set of all maximal weights of $V(\Lambda)$ by $\max(\Lambda)$. Hence, $\max(\Lambda) \cap P^+$ is the set of all maximal dominant weights of $V(\Lambda)$.

For $1 \leq \ell \leq \lfloor \frac{n}{2} \rfloor$, let $\gamma_\ell = \ell\alpha_0 + (\ell-1)\alpha_1 + (\ell-2)\alpha_2 + \cdots + \alpha_{\ell-1} + \alpha_{n-\ell+1} + \cdots + (\ell-2)\alpha_{n-2} + (\ell-1)\alpha_{n-1}$. It was shown in [5] that $\max(2\Lambda_0) \cap P^+ = \{2\Lambda_0 - \gamma_\ell \mid 1 \leq \ell \leq \lfloor \frac{n}{2} \rfloor\}$. Indeed, $\{k\Lambda_0 - \gamma_\ell \mid 1 \leq \ell \leq \lfloor \frac{n}{2} \rfloor\} \subseteq \max(k\Lambda_0) \cap P^+$ for $k \geq 2$ (see [1]). The following Theorem was proved in [1].

Theorem 3.1. [1] *For $k \geq 2$, $1 \leq \ell \leq \lfloor \frac{n}{2} \rfloor$, we have $mult_{k\Lambda_0}(k\Lambda_0 - \gamma_\ell) = |\mathcal{T}_\ell^k|$.*

Hence the following result is a consequence of Theorem 2.7 and Theorem 3.1.

Corollary 3.2. *For $k \geq 2$, $1 \leq \ell \leq \lfloor \frac{n}{2} \rfloor$, we have* $mult_{k\Lambda_0}(k\Lambda_0 - \gamma_\ell) = \sum_{\lambda \vdash \ell, h(\lambda) \leq k} (f^\lambda)^2.$

We denote a permutation of $\{1, 2, \ldots, \ell\}$ by a sequence $w = w_1 w_2 \ldots w_\ell$ if $1 \mapsto w_1, 2 \mapsto w_2, \ldots, \ell \mapsto w_\ell$. If a decreasing subsequence of w has m terms, we say it is a decreasing subsequence of length m. A $(k+1)k \ldots 21$-avoiding permutation is a permutation w with no decreasing subsequence of length $k+1$. For example, $w = 26873415$ has longest decreasing subsequence of length 4 (namely, 8731 and 8741) and so w is a 54321-avoiding permutation. We have the following result as a consequence of ([4], Corollary 7.23.12) and Theorem 3.1 which proves Conjecture 4.13 in [1].

Corollary 3.3. *For* $k \geq 2$, $1 \leq \ell \leq \lfloor \frac{n}{2} \rfloor$, *we have* $mult_{k\Lambda_0}(k\Lambda_0 - \gamma_\ell) =$ *the number of* $(k+1)k\cdots 21$-*avoiding permutations of* ℓ.

Example 3.4. Consider maximal dominant weight $4\Lambda_0 - \gamma_5$ of $V(4\Lambda_0)$. By Corollary 3.2, the multiplicity of $4\Lambda_0 - \gamma_5$ is equal to $\displaystyle\sum_{\lambda \vdash 5, h(\lambda) \leq 4} (f^\lambda)^2$. The partitions of 5 with height less than or equal to 4 are (5), (4,1), (3,2), (3,1,1),(2,2,1), and (2,1,1,1). In Table 1, we calculate f^λ for each of these partitions using the hook length formula. Thus, $mult_{4\Lambda_0}(4\Lambda_0 - \gamma_5) = 1^2 + 4^2 + 5^2 + 6^2 + 5^2 + 4^2 = 119$.

TABLE 1. Values of f^λ for $\lambda \vdash 5$, $ht(\lambda) \leq 4$

λ	f^λ
	$\frac{5!}{5\cdot 4\cdot 3\cdot 2\cdot 1} = 1$
	$\frac{5!}{5\cdot 3\cdot 2\cdot 1\cdot 1} = 4$
	$\frac{5!}{4\cdot 3\cdot 1\cdot 2\cdot 1} = 5$
	$\frac{5!}{5\cdot 2\cdot 1\cdot 2\cdot 1} = 6$
	$\frac{5!}{4\cdot 2\cdot 3\cdot 1\cdot 1} = 5$
	$\frac{5!}{5\cdot 1\cdot 3\cdot 2\cdot 1} = 4$

REFERENCES

[1] Jayne, R.L., Misra, K.C.: On multiplicities of maximal weights of $\widehat{sl}(n)$-modules, Algebr. Represent. Th. 17, 1303-1321 (2014)

[2] Kac, V.G.: Infinite-dimensional Lie algebras. Third edition. Cambridge University Press, New York (1990)

[3] Schensted, C.: Longest increasing and decreasing subsequences, Canadian J. Math. 13, 179-191 (1961)

[4] Stanley, R.P.: Enumerative Combinatorics. Vol. 2. Cambridge University Press, New York (1999)

[5] Tsuchioka, S.: Catalan numbers and level 2 weight structures of $A^{(1)}_{p-1}$, RIMS Kŏkyŭroku Bessatsu. B11, 145-154 (2009)

Hampden-Sydney College, Hampden-Sydney, VA 23943
E-mail address: `rjayne@hsc.edu`

Department of Mathematics, North Carolina State University, Raleigh, NC 27695-8205
E-mail address: `misra@ncsu.edu`